

Ginsparg-Wilson relation with $R = (a\gamma_5 D)^{2k}$

Ting-Wai Chiu^{a *}

^aDepartment of Physics, National Taiwan University, Taipei, Taiwan 106, Republic of China.

The Ginsparg-Wilson relation $D\gamma_5 + \gamma_5 D = 2aDR\gamma_5 D$ with $R = (a\gamma_5 D)^{2k}$ is discussed. An explicit realization of D is constructed. It is shown that this sequence of topologically-proper lattice Dirac operators tend to a nonlocal operator in the limit $k \rightarrow \infty$. This suggests that the locality of a lattice Dirac operator is irrelevant to its index.

1. INTRODUCTION

In general, the Ginsparg-Wilson relation [1] for a lattice Dirac operator D can be written as

$$D\gamma_5 f(D) + g(D)\gamma_5 D = 0, \quad (2)$$

where f and g are analytic functions. Then the fermionic action $\mathcal{A}_f = \bar{\psi} D \psi$ is invariant under the generalized chiral transformation

$$\psi \rightarrow \exp[\gamma_5 \theta f(D)] \psi, \quad (2)$$

$$\bar{\psi} \rightarrow \bar{\psi} \exp[\theta g(D)\gamma_5], \quad (3)$$

where θ is a global parameter. In particular, for $f(D) = \mathbf{1} - aRD$ and $g(D) = \mathbf{1} - aDR$, (1) becomes

$$D\gamma_5 + \gamma_5 D = aD(R\gamma_5 + \gamma_5 R)D, \quad (4)$$

where R is any operator. Since $(R\gamma_5 + \gamma_5 R)$ commutes with γ_5 , without loss, we can set R to commute with γ_5 ,

$$R\gamma_5 = \gamma_5 R. \quad (5)$$

Then (4) becomes the usual form of the GW relation

$$D\gamma_5 + \gamma_5 D = 2aDR\gamma_5 D. \quad (6)$$

In the continuum, the massless Dirac operator $\mathcal{D} = \gamma_\mu(\partial_\mu + iA_\mu)$ is chirally symmetric and anti-hermitian, i.e.,

$$D\gamma_5 + \gamma_5 D = 0, \quad (7)$$

$$\mathcal{D}^\dagger = -\mathcal{D}, \quad (8)$$

*This work was supported by the National Science Council, R.O.C. under the grant numbers NSC89-2112-M002-017 and NSC89-2112-M002-079

which imply that $\mathcal{D}$ is γ_5 -hermitian, i.e., $\mathcal{D}^\dagger = \gamma_5 \mathcal{D} \gamma_5$.

On the lattice, a lattice Dirac operator D cannot satisfy (7) or (8) without losing its other essential properties. However, we can still require D to satisfy the γ_5 -hermiticity,

$$D^\dagger = \gamma_5 D \gamma_5. \quad (9)$$

Multiplying γ_5 on both sides of (6), and using (9), we obtain $D^\dagger + D = 2aD^\dagger R D$, which gives $R^\dagger = R$.

Thus we conclude that, in general, R is a Hermitian operator commuting with γ_5 . From (6), the GW fermion propagator (in the trivial sector) satisfies $D^{-1}\gamma_5 + \gamma_5 D^{-1} = 2aR\gamma_5$, which immediately suggests that R should be as local as possible, i.e., $R \simeq \mathbf{1}$. Otherwise, it may cause additive mass renormalization to D^{-1} , which of course is undesirable. In fact, it has been demonstrated that the most optimal R is $R = r\mathbf{1}$ with some constant r [2].

Nevertheless, theoretically, it is rather interesting to consider R as a function of D , and to see how D behaves with respect to R . It turns out that, in general, the independent variable in the functional form of R must be $(a\gamma_5 D)^2$, if D is normal and γ_5 -hermitian. In particular, for $R(t) = t^n$, $n = 0, 1, \dots$, the sequence of topologically-proper D (27) tend to a nonlocal Dirac operator in the limit $n \rightarrow \infty$. This suggests that the locality of a lattice Dirac is irrelevant to its index.

First, recall that for any D which is normal ($DD^\dagger = D^\dagger D$) and γ_5 -hermitian, its eigensystem ($D\phi_s = \lambda_s\phi_s$) has the following properties [3]:

- (i) eigenvalues are either real or come in complex conjugate pairs.
- (ii) the chirality ($\chi_s = \phi_s^\dagger \gamma_5 \phi_s$) of any complex eigenmode is zero.
- (iii) each real eigenmode has a definite chirality.
- (iv) sum of the chirality of all real eigenmodes is zero (chirality sum rule).

Assume that the eigenvalues of D lying on a simple closed contour in the complex plane. Then (i) implies that the real eigenvalues of D can only occur at two different points, say at zero and a^{-1} . Then the chirality sum rule reads

$$\sum_{\lambda_s=0, a^{-1}} \phi_s^\dagger \gamma_5 \phi_s = 0, \quad (10)$$

which immediately gives

$$n_+ - n_- + N_+ - N_- = 0. \quad (11)$$

where $n_\pm$ denotes the number of zero modes of $\pm$ chirality, and $N_\pm$ the number of nonzero real (a^{-1}) eigenmodes of $\pm$ chirality. Then we immediately see that any zero mode must be accompanied by a real (a^{-1}) eigenmode with opposite chirality, and the index of D is

$$\text{index}(D) \equiv n_- - n_+ = -(N_- - N_+). \quad (12)$$

It should be emphasized that the chiral properties (ii), (iii) and the chirality sum rule (11) hold for any normal D satisfying the γ_5 -hermiticity, as shown in Ref. [3]. However, in nontrivial gauge backgrounds, whether D possesses any zero modes or not relies on the topological characteristics [11] of D , which cannot be guaranteed by the conditions such as the locality, free of species doublings, correct continuum behavior, γ_5 -hermiticity and the GW relation.

Therefore, we must require that D is normal and γ_5 -hermitian,

$$D^\dagger D = D D^\dagger, \quad D^\dagger = \gamma_5 D \gamma_5.$$

These two equations immediately give

$$\gamma_5 (\gamma_5 D)^2 = (\gamma_5 D)^2 \gamma_5. \quad (13)$$

Thus we have found an example of $R = (a\gamma_5 D)^2$ which is Hermitian and commutes with γ_5 . In

general, R can be any analytic function of $(a\gamma_5 D)^2$, i.e.,

$$R = f[(a\gamma_5 D)^2], \quad f : \text{analytic function}. \quad (14)$$

In particular, for $f(x) = x^n$, it becomes

$$R = (a\gamma_5 D)^{2k}, \quad k = 0, 1, 2, \dots \quad (15)$$

Substituting (15) into the GW relation (6), we obtain

$$\gamma_5 D + D \gamma_5 = 2aD(a\gamma_5 D)^{2k} \gamma_5 D, \quad (16)$$

which is equivalent to Fujikawa's proposal [4]

$$\gamma_5 (\gamma_5 D) + (\gamma_5 D) \gamma_5 = 2a^{2k+1} (\gamma_5 D)^{2k+2}. \quad (17)$$

2. A CONSTRUCTION OF D

So far, the only viable way to construct a topologically-proper lattice Dirac operator is the Overlap [5]

$$D = \frac{1}{2a}(\mathbf{I} + \gamma_5 \epsilon), \quad \epsilon^2 = \mathbf{I} \quad (18)$$

which satisfies the GW relation (6) with $R = 1$. Since D (18) is normal and γ_5 -hermitian, its eigenmodes satisfy the chiral properties (i)-(iv). There are many different ways to implement the Hermitian ϵ in (18). However, it is required to be able to capture the topology of the gauge background. That means, one-half of the difference of the numbers ($h_\pm$) of positive ($+1$) and negative (-1) eigenvalues of ϵ is equal to the background topological charge Q ,

$$\sum_x \text{tr}(a\gamma_5 D(x, x)) = \frac{1}{2} \sum_x \text{tr}(\epsilon) = \frac{1}{2}(h_+ - h_-) = Q$$

where tr denotes the trace over the Dirac and color space. Otherwise, the axial anomaly of D cannot agree with the topological charge density in a nontrivial gauge background.

An explicit realization of ϵ in (18) is the Neuberger-Dirac operator [6] with

$$\epsilon = \frac{H_w}{\sqrt{H_w^2}} \quad (19)$$

where

$$H_w = \gamma_5 (\gamma_\mu t_\mu + W - m_0 a^{-1}), \quad 0 < m_0 < 2 \quad (20)$$

$$t_\mu(x, y) = \frac{1}{2a} [U_\mu(x)\delta_{x+\hat{\mu}, y} - U_\mu^\dagger(y)\delta_{x-\hat{\mu}, y}] \quad (21)$$

$$W(x, y) = \frac{1}{2a} \sum_\mu [2\delta_{x, y} - U_\mu(x)\delta_{x+\hat{\mu}, y} - U_\mu^\dagger(y)\delta_{x-\hat{\mu}, y}]$$

To generalize this construction for $k > 0$, we multiply both sides of the GW relation (6) by R and redefine $D' = RD$, then we have

$$D'\gamma_5 + \gamma_5 D' = 2D'\gamma_5 D' . \quad (22)$$

It can be shown that D' is γ_5 -hermitian. Now (22) is in the same form of the GW relation with $R = 1$. Thus, one can construct D' in the same way as the Overlap

$$D' = RD = \frac{1}{2a}(\mathbf{1} + \gamma_5 \epsilon), \quad \epsilon^2 = \mathbf{1}, \quad (23)$$

provided that a proper realization of ϵ can be obtained. Using (5), we obtain

$$D' = RD = a^{-1}\gamma_5(a\gamma_5 D)^{2k+1}, \quad (24)$$

which immediately yields

$$D = a^{-1}\gamma_5(a\gamma_5 D')^{1/(2k+1)}, \quad (25)$$

where the $(2k+1)$ -th real root of the Hermitian operator $(a\gamma_5 D')$ is assumed. Then (25) suggests that if the ϵ in (23) is expressed in terms of a Hermitian operator H , i.e., $\epsilon = \frac{H}{\sqrt{H^2}}$, then H is required to behave as $(\gamma_\mu \mathcal{D}_\mu)^{2k+1}$ in the continuum limit such that D can become $\gamma_\mu \mathcal{D}_\mu$ after taking the $(2k+1)$ -th root in (25). Thus, H must contain the term $(\gamma_\mu t_\mu)^{2k+1}$, where $\gamma_\mu t_\mu$ is the naive lattice fermion operator defined in (21). Then another term must be added in order to remove the species doublers in $(\gamma_\mu t_\mu)^{2k+1}$. So, we add the Wilson term to the $(2k+1)$ -th power, i.e., W^{2k+1} . Finally, a negative mass term $-\frac{1}{2}a^{-(2k+1)}$ is inserted such that ϵ is able to detect the topological charge Q of the gauge background, i.e.,

$$\frac{1}{2} \sum_x \text{tr} \left(\frac{H}{\sqrt{H^2}} \right) = Q. \quad (26)$$

Putting all these terms together, we have

$$H = \gamma_5 \left[(-1)^k (\gamma_\mu t_\mu)^{2k+1} + W^{2k+1} - \frac{a^{-(2k+1)}}{2} \right]$$

which, at $k = 0$, reduces to the H_w in Eq. (20). Then (25) can be rewritten as

$$D = a^{-1} \left(\frac{1}{2} \right)^{\frac{1}{(2k+1)}} \gamma_5 \left(\gamma_5 + \frac{H}{\sqrt{H^2}} \right)^{\frac{1}{(2k+1)}} \quad (27)$$

which agrees with Fujikawa's construction [4]. Note that (27) can also be written in the form $D = D_c(\mathbf{1} + aRD_c)^{-1}$, where D_c is dependent on the order k . The general properties of D have been derived in [7]. In particular, the index of D is independent of the order k , and the eigenvalues of D fall on a closed contour with their real parts bounded between zero and a^{-1} . In Fig. 1, the eigenvalues of D are plotted for $k = 0, 1, 2$ respectively, with the same background gauge field.

3. IN THE LIMIT $k \rightarrow \infty$

One of the salient features of the sequence of topologically-proper Dirac operators (27) is that the amount of chiral symmetry breaking (i.e., r.h.s. of (6)) decreases as the order k increases. However, at finite lattice spacing, the chiral symmetry breaking of D *cannot* be zero even in the limit $k \rightarrow \infty$, since D satisfies the GW relation (6). In the limit $k \rightarrow \infty$, the only possibility for D to break the chiral symmetry is that $D(p)$ becomes a piecewise continuous function in the Brillouin zone, with discontinuities somewhere at $|p| \simeq a^{-1}$, and $\gamma_5 D(p) + D(p)\gamma_5 = 0$ for $|p| < a^{-1}$. Since such a $D(p)$ is non-analytic at infinite number of p , the corresponding $D(x, y)$ must be non-local in the position space.

If $D(x, y)$ is nonlocal in the free fermion limit (i.e., $U_\mu(x) \rightarrow 1$), then it must be nonlocal in any smooth gauge background. So, it suffices to examine the non-analyticity of $D(p)$ in the free fermion limit.

In the free fermion limit, the GW Dirac operator (27) in momentum space can be written as

$$D(p) = D_0(p) + i\gamma_\mu D_\mu(p), \quad (28)$$

where

$$D_0(p) = a^{-1} \left[\frac{1}{2} \left(1 + \frac{u(p)}{N(p)} \right) \right]^{\frac{k+1}{2k+1}},$$

$$D_\mu(p) = D_0(p) \sqrt{\frac{1}{a^2 t^2(p)} \left(\frac{N(p) - u(p)}{N(p) + u(p)} \right)} \sin(p_\mu a),$$




Figure 1. The eigenvalues of D in a nontrivial gauge background for $k = 0, 1, 2$ respectively.

$$u(p) = a^{-(2k+1)} \left(\left[\sum_{\mu} (1 - \cos(p_{\mu}a)) \right]^{2k+1} - \frac{1}{2} \right)$$

$$t^2(p) = a^{-2} \sum_{\mu} \sin^2(p_{\mu}a) ,$$

$$N(p) = \sqrt{[t^2(p)]^{2k+1} + [u(p)]^2} .$$

At the zeroth order ($k = 0$), $D(p)$ is analytic in the Brillouin zone. However, in the limit $k \rightarrow \infty$, $D(p)$ tends to a nonanalytic function with discontinuities at each point on two concentric hypersurfaces inside the Brillouin zone [8]. In four dimensions, the inscribed hypersurface is specified by the equation

$$\sum_{\mu=1}^4 s_{\mu}^2 = 1 ,$$

while the circumscribing one by

$$6 + \sum_{\mu=1}^4 c_{\mu}(c_{\mu} - 4) + \sum_{\mu=1}^3 \sum_{\nu > \mu}^4 c_{\mu}c_{\nu} = 0 ,$$

where $c_{\mu} = \cos(p_{\mu}a)$ and $s_{\mu} = \sin(p_{\mu}a)$. This implies that $D(x, y)$ is nonlocal in the limit $k \rightarrow \infty$.

On the other hand, $D(x, y)$ is local at the zeroth order ($k = 0$), for gauge backgrounds which are sufficiently smooth at the scale of the lattice spacing [9,10]. Since the index of D (27) is independent of the order k , this suggests that the locality of a lattice Dirac operator is irrelevant to its index. In other words, if a GW Dirac operator which is γ_5 -hermitian, doubler-free, and has correct continuum behavior in the free fermion limit, *but* always has zero index [11], then the cause may *not* be due to its nonlocality.

To summarize, we have provided a nontrivial example of D (27), which is *topologically-proper* but tends to a *nonlocal* operator in the limit $k \rightarrow \infty$. Unlike the nonlocal and chirally-symmetric Dirac operator

$$D_c = a^{-1} \frac{\sqrt{H_w^2} + \gamma_5 H_w}{\sqrt{H_w^2} - \gamma_5 H_w} \quad (29)$$

which has poles (i.e., the nonzero real eigenmodes) for nontrivial background gauge fields (as a consequence of the chirality sum rule [3]), the present example (27) breaks the chiral symmetry according to the GW relation, thus its nonzero real (a^{-1}) eigenmodes are well-defined for any order k .

REFERENCES

1. P. Ginsparg and K. Wilson, Phys. Rev. D25 (1982) 2649.
2. T.W. Chiu, Phys. Lett. B474 (2000) 89.
3. T.W. Chiu, Phys. Rev. D58 (1998) 074511.
4. K. Fujikawa, Nucl. Phys. B589 (2000) 487.
5. R. Narayanan, H. Neuberger, Nucl. Phys. B443 (1995) 305.
6. H. Neuberger, Phys. Lett. B417 (1998) 141; Phys. Lett. B427 (1998) 353.
7. T.W. Chiu, Nucl. Phys. B588 (2000) 400.
8. T.W. Chiu, hep-lat/0008010.
9. P. Hernández, K. Jansen, M. Lüscher, Nucl. Phys. B552 (1999) 363.
10. H. Neuberger, Phys. Rev. D61 (2000) 085015.
11. T.W. Chiu, hep-lat/9911010.